


**Navane®**  
**thiothixene hydrochloride**  
**Intramuscular For Injection**  
**STERILE**

**DESCRIPTION**

Navane (thiothixene hydrochloride) is a thioxanthene derivative. Specifically, thiothixene is the *cis* isomer of N,N-dimethyl-9-[3-(4-methyl-1-piperazinyl)-propylidene] thioxanthene-2-sulfonamide.

The thioxanthenes differ from the phenothiazines by the replacement of nitrogen in the central ring with a carbon-linked side chain fixed in space in a rigid structural configuration. An N,N-dimethyl sulfonamide functional group is bonded to the thioxanthene nucleus.



thiothixene hydrochloride

Inert ingredients for the intramuscular for injection formulation are: water; mannitol.

**ACTIONS**

Navane is a psychotropic agent of the thioxanthene series. Navane possesses certain chemical and pharmacological similarities to the piperazine phenothiazines and differences from the aliphatic group of phenothiazines. Navane's mode of action has not been clearly established.

**INDICATIONS**

Navane is effective in the management of manifestations of psychotic disorders. Navane has not been evaluated in the management of behavioral complications in patients with mental retardation.

**CONTRAINDICATIONS**

Navane is contraindicated in patients with circulatory collapse, comatose states, central nervous system depression due to any cause, and blood dyscrasias. Navane is contraindicated in

individuals who have shown hypersensitivity to the drug. It is not known whether there is a cross sensitivity between the thioxanthenes and the phenothiazine derivatives, but this possibility should be considered.

### WARNINGS

**Tardive Dyskinesia**—Tardive dyskinesia, a syndrome consisting of potentially irreversible, involuntary, dyskinetic movements may develop in patients treated with neuroleptic (antipsychotic) drugs. Although the prevalence of the syndrome appears to be highest among the elderly, especially elderly women, it is impossible to rely upon prevalence estimates to predict, at the inception of neuroleptic treatment, which patients are likely to develop the syndrome. Whether neuroleptic drug products differ in their potential to cause tardive dyskinesia is unknown.

Both the risk of developing the syndrome and the likelihood that it will become irreversible are believed to increase as the duration of treatment and the total cumulative dose of neuroleptic drugs administered to the patient increase. However, the syndrome can develop, although much less commonly, after relatively brief treatment periods at low doses.

There is no known treatment for established cases of tardive dyskinesia, although the syndrome may remit, partially or completely, if neuroleptic treatment is withdrawn. Neuroleptic treatment, itself, however, may suppress (or partially suppress) the signs and symptoms of the syndrome and thereby may possibly mask the underlying disease process. The effect that symptomatic suppression has upon the long-term course of the syndrome is unknown.

Given these considerations, neuroleptics should be prescribed in a manner that is most likely to minimize the occurrence of tardive dyskinesia. Chronic neuroleptic treatment should generally be reserved for patients who suffer from a chronic illness that, 1) is known to respond to neuroleptic drugs, and, 2) for whom alternative, equally effective, but potentially less harmful treatments are *not* available or appropriate. In patients who do require chronic treatment, the smallest dose and the shortest duration of treatment producing a satisfactory clinical response should be sought. The need for continued treatment should be reassessed periodically.

If signs and symptoms of tardive dyskinesia appear in a patient on neuroleptics, drug discontinuation should be considered. However, some patients may require treatment despite the presence of the syndrome.

(For further information about the description of tardive dyskinesia and its clinical detection, please refer to “Information for Patients” in the PRECAUTIONS section, and to the ADVERSE REACTIONS section.)

**Neuroleptic Malignant Syndrome (NMS)**—A potentially fatal symptom complex sometimes referred to as Neuroleptic Malignant Syndrome (NMS) has been reported in association with antipsychotic drugs. Clinical manifestations of NMS are hyperpyrexia, muscle rigidity, altered mental status and evidence of autonomic instability (irregular pulse or blood pressure, tachycardia, diaphoresis, and cardiac dysrhythmias).

The diagnostic evaluation of patients with this syndrome is complicated. In arriving at a diagnosis, it is important to identify cases where the clinical presentation includes both serious medical illness (e.g., pneumonia, systemic infection, etc.) and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever and primary central nervous system (CNS) pathology.

The management of NMS should include 1) immediate discontinuation of antipsychotic drugs and other drugs not essential to concurrent therapy, 2) intensive symptomatic treatment and medical monitoring, and 3) treatment of any concomitant serious medical problems for which specific treatments are available. There is no general agreement about specific pharmacological treatment regimens for uncomplicated NMS.

If a patient requires antipsychotic drug treatment after recovery from NMS, the potential reintroduction of drug therapy should be carefully considered. The patient should be carefully monitored, since recurrences of NMS have been reported.

**Usage in Pregnancy**—Safe use of Navane during pregnancy has not been established. Therefore, this drug should be given to pregnant patients only when, in the judgment of the physician, the expected benefits from treatment exceed the possible risks to mother and fetus. Animal reproductive studies and clinical experience to date have not demonstrated any teratogenic effects.

In the animal reproduction studies with Navane, there was some decrease in conception rate and litter size, and an increase in resorption rate in rats and rabbits, changes which have been similarly reported with other psychotropic agents. After repeated oral administration of Navane to rats (5 to 15 mg/kg/day), rabbits (3 to 50 mg/kg/day), and monkeys (1 to 3 mg/kg/day) before and during gestation, no teratogenic effects were seen. (See Precautions.)

**Usage in Children**—The use of Navane in children under 12 years of age is not recommended because safety and efficacy in the pediatric age group have not been established.

As is true with many CNS drugs, Navane may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery, especially during the first few days of therapy. Therefore, the patient should be cautioned accordingly.

As in the case of other CNS-acting drugs, patients receiving Navane should be cautioned about the possible additive effects (which may include hypotension) with CNS depressants and with alcohol.

### PRECAUTIONS

**General:** An antiemetic effect was observed in animal studies with Navane (thiothixene hydrochloride); since this effect may also occur in man, it is possible that Navane may mask signs of overdosage of toxic drugs and may obscure conditions such as intestinal obstruction and brain tumor.

In consideration of the known capability of Navane and certain other psychotropic drugs to precipitate convulsions, extreme caution should be used in patients with a history of convulsive disorders, or those in a state of alcohol withdrawal since it may lower the convulsive threshold. Although Navane potentiates the actions of the barbiturates, the dosage of the anticonvulsant therapy should not be reduced when Navane is administered concurrently.

Caution as well as careful adjustment of the dosage is indicated when Navane is used in conjunction with other CNS depressants other than anticonvulsant drugs.

Though exhibiting rather weak anticholinergic properties, Navane should be used with caution in patients who are known or suspected to have glaucoma, or who might be exposed to extreme heat, or who are receiving atropine or related drugs.

Use with caution in patients with cardiovascular disease.

Also, careful observation should be made for pigmentary retinopathy, and lenticular pigmentation (fine lenticular pigmentation has been noted in a small number of patients treated with Navane for prolonged periods). Blood dyscrasias (agranulocytosis, pancytopenia, thrombocytopenic purpura), and liver damage (jaundice, biliary stasis), have been reported with related drugs.

Undue exposure to sunlight should be avoided. Photosensitive reactions have been reported in patients on Navane.

As with all intramuscular preparations, Navane Intramuscular For Injection should be injected well within the body of a relatively large muscle. The preferred sites are the upper outer quadrant of the buttock (i.e., gluteus maximus) and the mid-lateral thigh.

The deltoid area should be used only if well developed such as in certain adults and older children, and then only with caution to avoid radial nerve injury. Intramuscular injections should not be made into the lower and mid-thirds of the upper arm. As with all intramuscular injections, aspiration is necessary to help avoid inadvertent injection into a blood vessel.

Neuroleptic drugs elevate prolactin levels; the elevation persists during chronic administration. Tissue culture experiments indicate that approximately one-third of human breast cancers are prolactin dependent *in vitro*, a factor of potential importance if the prescription of these drugs is

contemplated in a patient with a previously detected breast cancer. Although disturbances such as galactorrhea, amenorrhea, gynecomastia, and impotence have been reported, the clinical significance of elevated serum prolactin levels is unknown for most patients. An increase in mammary neoplasms has been found in rodents after chronic administration of neuroleptic drugs. Neither clinical studies nor epidemiologic studies conducted to date, however, have shown an association between chronic administration of these drugs and mammary tumorigenesis; the available evidence is considered too limited to be conclusive at this time.

**Information for Patients:** Given the likelihood that some patients exposed chronically to neuroleptics will develop tardive dyskinesia, it is advised that all patients in whom chronic use is contemplated be given, if possible, full information about this risk. The decision to inform patients and/or their guardians must obviously take into account the clinical circumstances and the competency of the patient to understand the information provided.

### **ADVERSE REACTIONS**

NOTE: Not all of the following adverse reactions have been reported with Navane. However, since Navane has certain chemical and pharmacologic similarities to the phenothiazines, all of the known side effects and toxicity associated with phenothiazine therapy should be borne in mind when Navane is used.

**Cardiovascular Effects:** Tachycardia, hypotension, lightheadedness, and syncope. In the event hypotension occurs, epinephrine should not be used as a pressor agent since a paradoxical further lowering of blood pressure may result. Nonspecific EKG changes have been observed in some patients receiving Navane. These changes are usually reversible and frequently disappear on continued Navane therapy. The clinical significance of these changes is not known.

**CNS Effects:** Drowsiness, usually mild, may occur although it usually subsides with continuation of Navane therapy. The incidence of sedation appears similar to that of the piperazine group of phenothiazines, but less than that of certain aliphatic phenothiazines. Restlessness, agitation and insomnia have been noted with Navane. Seizures and paradoxical exacerbation of psychotic symptoms have occurred with Navane infrequently.

Hyperreflexia has been reported in infants delivered from mothers having received structurally related drugs.

In addition, phenothiazine derivatives have been associated with cerebral edema and cerebrospinal fluid abnormalities.

### **Extrapyramidal Symptoms**

Extrapyramidal symptoms, such as pseudo-parkinsonism, akathisia, and dystonia have been reported (see Dystonia, Class effect). Management of these extrapyramidal symptoms depends upon the type and severity. Rapid relief of acute symptoms may require the use of an injectable antiparkinson agent. More slowly emerging symptoms may be managed by reducing the dosage of Navane and/or administering an oral antiparkinson agent.

## **Dystonia**

***Class effect:*** Symptoms of dystonia, prolonged abnormal contractions of muscle groups, may occur in susceptible individuals during the first few days of treatment. Dystonic symptoms include: spasm of the neck muscles, sometimes progressing to tightness of the throat, swallowing difficulty, difficulty breathing, and/or protrusion of the tongue. While these symptoms can occur at low doses, they occur more frequently and with greater severity with high potency and at higher doses of first generation antipsychotic drugs. An elevated risk of acute dystonia is observed in males and younger age groups.

**Persistent Tardive Dyskinesia:** As with all antipsychotic agents, tardive dyskinesia may appear in some patients on long-term therapy or may occur after drug therapy has been discontinued. The syndrome is characterized by rhythmical involuntary movements of the tongue, face, mouth or jaw (e.g., protrusion of tongue, puffing of cheeks, puckering of mouth, chewing movements). Sometimes these may be accompanied by involuntary movements of extremities.

Since early detection of tardive dyskinesia is important, patients should be monitored on an ongoing basis. It has been reported that fine vermicular movement of the tongue may be an early sign of the syndrome. If this or any other presentation of the syndrome is observed, the clinician should consider possible discontinuation of neuroleptic medication. (See WARNINGS section.)

**Hepatic Effects:** Elevations of serum transaminase and alkaline phosphatase, usually transient, have been infrequently observed in some patients. No clinically confirmed cases of jaundice attributable to Navane (thiothixene hydrochloride) have been reported.

**Hematologic Effects:** As is true with certain other psychotropic drugs, leukopenia and leucocytosis, which are usually transient, can occur occasionally with Navane. Other antipsychotic drugs have been associated with agranulocytosis, eosinophilia, hemolytic anemia, thrombocytopenia and pancytopenia.

**Allergic Reactions:** Rash, pruritus, urticaria, and rare cases of anaphylaxis have been reported with Navane. Undue exposure to sunlight should be avoided. Although not experienced with Navane, exfoliative dermatitis, contact dermatitis (in nursing personnel), have been reported with certain phenothiazines.

**Endocrine Disorders:** Lactation, moderate breast enlargement and amenorrhea have occurred in a small percentage of females receiving Navane. If persistent, this may necessitate a reduction in dosage or the discontinuation of therapy. Phenothiazines have been associated with false positive pregnancy tests, gynecomastia, hypoglycemia, hyperglycemia, and glycosuria.

**Autonomic Effects:** Dry mouth, blurred vision, nasal congestion, constipation, increased sweating, increased salivation, and impotence have occurred infrequently with Navane therapy. Phenothiazines have been associated with miosis, mydriasis, and adynamic ileus.

Other Adverse Reactions: Hyperpyrexia, anorexia, nausea, vomiting, diarrhea, increase in appetite and weight, weakness or fatigue, polydipsia and peripheral edema.

Although not reported with Navane, evidence indicates there is a relationship between phenothiazine therapy and the occurrence of a systemic lupus erythematosus-like syndrome.

Neuroleptic Malignant Syndrome (NMS): Please refer to the text regarding NMS in the WARNINGS section.

NOTE: Sudden deaths have occasionally been reported in patients who have received certain phenothiazine derivatives. In some cases the cause of death was apparently cardiac arrest or asphyxia due to failure of the cough reflex. In others, the cause could not be determined nor could it be established that death was due to phenothiazine administration.

## **DOSAGE AND ADMINISTRATION**

### **Preparation**

Navane (thiothixene hydrochloride) Intramuscular For Injection must be reconstituted with 2.2 mL of sterile water for injection.

### **For Intramuscular Use Only**

Dosage of Navane should be individually adjusted depending on the chronicity and severity of the condition. In general, small doses should be used initially and gradually increased to the optimal effective level, based on patient response.

Usage in children under 12 years of age is not recommended.

Where more rapid control and treatment of acute behavior is desirable, the intramuscular form of Navane may be indicated. It is also of benefit where the very nature of the patient's symptomatology, whether acute or chronic, renders oral administration impractical or even impossible.

For treatment of acute symptomatology or in patients unable or unwilling to take oral medication, the usual dose is 4 mg of Navane Intramuscular For Injection administered 2 to 4 times daily. Dosage may be increased or decreased depending on response. Most patients are controlled on a total daily dosage of 16 to 20 mg. The maximum recommended dosage is 30 mg/day. An oral form should supplant the injectable form as soon as possible. It may be necessary to adjust the dosage when changing from the intramuscular to oral dosage forms. Dosage recommendations for Navane Capsules and Concentrate can be found in the Navane oral package insert.

## **OVERDOSAGE**

Manifestations include muscular twitching, drowsiness, and dizziness. Symptoms of gross overdosage may include CNS depression, rigidity, weakness, torticollis, tremor, salivation, dysphagia, hypotension, disturbances of gait, or coma.

Treatment: Essentially symptomatic and supportive. Keep patient under careful observation and maintain an open airway, since involvement of the extrapyramidal system may produce dysphagia and respiratory difficulty in severe overdosage. If hypotension occurs, the standard measures for managing circulatory shock should be used (I.V. fluids and/or vasoconstrictors).

If a vasoconstrictor is needed, levarterenol and phenylephrine are the most suitable drugs. Other pressor agents, including epinephrine, are not recommended, since phenothiazine derivatives may reverse the usual pressor elevating action of these agents and cause further lowering of blood pressure.

If CNS depression is marked, symptomatic treatment is indicated. Extrapyramidal symptoms may be treated with antiparkinson drugs.

There are no data on the use of peritoneal or hemodialysis, but they are known to be of little value in phenothiazine intoxication.

#### **HOW SUPPLIED**

Navane (thiothixene hydrochloride) Intramuscular For Injection is available in amber glass vials in packages of 10 vials (NDC 0049-5765-83). When reconstituted with 2.2 mL of STERILE WATER FOR INJECTION, each mL contains thiothixene hydrochloride equivalent to 5 mg of thiothixene, and 59.6 mg of mannitol. The reconstituted solution of Navane Intramuscular For Injection may be stored for 48 hours at room temperature before discarding.

**Rx only**



*Distributed by*

---

**Roerig**

Division of Pfizer Inc, NY, NY 10017

LAB-0252-3.0  
Revised January 2008